# REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information, Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE  2. October 2003	0	ND DATES COVERED /99 — 06/30/03)
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS
Medical Ultrasound Technology Rese	earch and Development a	t the University of	
Washington Center for Industrial and Medical Ultrasound			N00014-99-1-0982
6. AUTHOR(S)			
Lawrence A. Crum		•	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS	C(ES)		8. PERFORMING ORGANIZATION
Applied Physics Laboratory			REPORT NUMBER
University of Washington			
1013 NE 40th St.			
Seattle, WA 98105-6698			
O ODDANOCINO A CONITODINO A CENCY MANACIO AND AD	DDE66/E6)		10. SPONSORING/MONITORING
9. SPONSORING/MONITORING AGENCY NAME(S) AND AD Office of Naval Research	DHE99(E9)		AGENCY REPORT NUMBER
800 N. Quincy St.			1
Arlington, VA 22217-5660			1
Attn: Dr. Michael Given, Code 3-	41	I	

11. SUPPLEMENTARY NOTES

20031106 085

13. ABSTRACT (Maximum 200 words)

12a, DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release

This grant provided support to expand the scientific program and infrastructure at the University of Washington's Center for Industrial and Medical Ultrasound (CIMU). The many disparate facilities and technical capabilities available to CIMU staff and students were integrated and enhanced to provide a world-class, advanced research center for bioengineering development and graduate education in high-intensity, focused ultrasound (HIFU). This included leveraging of overall research in understanding of acoustic hemostasis and improvement of research tools. Significant progress was made in developing a highly automated transducer characterization capability. Features include acoustic field mapping by scanned hydrophones, acoustic power efficiency measurement, electric and acoustic impedance measurement, medical ultrasound imaging and RF data collection, optical methods of acoustic field mapping (Schlieren apparatus), and for each system a convenient software interface and user manual. Specialized fixtures and instruments were developed to measure HIFU dose response in vitro, an essential component of developing reliable and realistic computer models for HIFU simulation and treatment planning. Single-element HIFU transducers were refined with liquid, gel, and solid coupling media and other improvements to be more powerful, convenient, and robust. These technological enhancements have enabled the development of HIFU arrays and image-guided ultrasound systems for greater flexibility and control in clinical treatment protocol. Lastly, significant bioacoustic model development was undertaken for use in practical engineering design as well as for fundamental research of the mechanisms of ultrasound therapy.

14. SUBJECT TERMS	15. NUMBER OF PAGES 17		
acoustic hemostasis, bioacoustic (HIFU), ultrasound	16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)

# FINAL TECHNICAL REPORT FOR ONR GRANT N00014-99-1-0982 "MEDICAL ULTRASOUND TECHNOLOGY RESEARCH AND DEVELOPMENT AT THE UNIVERSITY OF WASHINGTON CENTER FOR INDUSTRIAL AND MEDICAL ULTRASOUND"

Lawrence A. Crum, Principal Investigator

# I. Background

This grant supported a research effort to understand more fully the basic science issues that comprise High Intensity Focused Ultrasound (HIFU) technology, and to build the basis for more expansive development of this general technology. Specific areas on interest included the use of HIFU in the treatment of hemorrhagic trauma and related pathological conditions, especially in organs that are difficult to treat using conventional medical and surgical techniques. Direct applications include combat casualty care, as well as many civilian uses in non-invasive or minimally invasive trauma management, bloodless surgery, and ultrasound-mediated drug therapy. Research efforts also explored imaging and simulation techniques associated with treatment, targeting and monitoring the effectiveness of HIFU therapy. A significant portion of the grant was directed toward renovation of laboratory spaces to enable the research efforts described above. The period of performance of the grant was from 1 July1999 to 30 June 2003.

# II. Accomplishments

# A. Executive Summary

This grant provided support to expand the scientific program and infrastructure at the University of Washington (UW) Center for Industrial and Medical Ultrasound (CIMU). The broad objectives falling within this scope included:

- To understand more fully the physical and biological mechanisms that give rise to acoustic hemostasis and HIFU surgical lesion production.
- To develop techniques for the utilization of gas-filled ultrasound contrast agents in the detection and localization of bleeding.
- To develop and validate theoretical models of acoustic hemostasis and HIFU surgical lesion production.
- To develop techniques for the real-time monitoring of HIFU-induced lesions in tissue.
- To develop additional laboratory transducer designs that produce acoustic hemostasis and HIFU lesions, and to extend these laboratory designs toward engineering prototypes.
- To understand more fully and to optimize the physical and biological mechanisms that give rise to ultrasound-mediated drug delivery.

During the performance period the many disparate facilities and technical capabilities available to CIMU staff and students were integrated and enhanced to provide a world-class, advanced research center for HIFU bioengineering development and graduate education. Program funds were used to leverage overall research efforts in understanding of acoustic hemostasis and improve research tools. Significant progress was made in developing a highly automated transducer characterization capability, including providing dedicated instrumentation and facilities for this purpose. Features include acoustic field mapping by scanned hydrophones, acoustic power efficiency measurement, electric and acoustic impedance measurement, acoustic and material parameter measurement, medical ultrasound imaging and RF data collection, optical methods of acoustic field mapping (Schlieren apparatus), and for each system a convenient software interface and user manual. Specialized fixtures and instruments were developed to measure HIFU dose response in vitro, an essential component of developing reliable and realistic computer models for HIFU simulation and treatment planning. Single element HIFU transducers were refined with liquid, gel, and solid coupling media and other improvements to be more powerful, convenient, and robust. These technological enhancements have enabled the development of HIFU arrays and image-guided ultrasound systems for greater flexibility and control in clinical HIFU treatment protocol. Lastly, significant bioacoustic model development was undertaken for use in practical engineering design as well as for fundamental research of the mechanisms of ultrasound therapy.

## B. R&D Facilities Improvements


A significant portion of this grant was directed toward renovation of laboratories to support research efforts at the newly created Center for Industrial and Medical Ultrasound. Space was acquired in the University of Washington Old Fisheries Center (OFC), conveniently located near the UW's Medical Center and Health Sciences buildings, but facilities improvements were urgently needed. Working in careful coordination with facilities crews from both the Applied Physics Laboratory and Main Campus, the best possible use was made of renovation funds provided under this grant.

One of the major benefits of the laboratory renovation has been to create good quality laboratory facilities that are proximal to the clinical facilities i.e. the UW Medical School and Medical Center. This location allows clinicians to be able to interact with the laboratories and still be closely available to their clinical duties. This becomes increasingly important as diagnostic and therapeutic ultrasound projects involve more clinical input. It is also more convenient for the subjects/patients. The proximity of the location to the Dept. of Radiology is particularly important for projects such as vascular ultrasound where input is needed on imaging and image guided therapy. The ultrasound patient studies laboratory (OFC room 119) provides specially-needed facilities for obtaining 3D images, eg. a non-magnetic environment for image registration.

Other highlights of the renovation efforts are described and illustrated in the following paragraphs.

## OFC Room 102N: Joint Imaging Laboratory


This laboratory was the most significantly in need of repair, as it contained unidentified chemicals, discarded nonfunctional equipment, and peeling lead-based wall and fixture paint. Lead-based paint abatement was completed, new paint applied, and the rusted countertops were sanded and buffed; before and after photographs are shown in Figure 1. This lab now serves as the joint imaging lab, used by multiple investigators working on new ultrasound imaging and detection algorithms, and systems integration for image-guided therapy.




#### OFC Room 112: Ultrasound Biophysics Laboratory

This laboratory was completely remodeled to serve as an ultrasound bioeffects laboratory for both *in vitro* and *in vivo* experimentation. It had been a radiation biophysics lab with extremely limited peripheral bench surfaces, no work surface in the center of the room, wornout wooden cabinetry, and very low quality water supply. The original walls contained asbestos, which required significant abatement efforts. The laboratory now features modern steel cabinetry, spacious bench areas, including a sizable central island affording unrestricted access to large and complex experimental setups (or animals) from all sides, and providing a great deal of flexibility with respect to equipment configuration. All surfaces are clad in

impermeable materials and can be chemically disinfected to remove either potentially biohazardous contaminants or to prepare surfaces for animal research. Abundant 120 V (and one 220 V) power drops were installed overhead to permit the use of a large number electronic instruments without encumbering the work surfaces with power cords. The lab now has two spatially separate sink areas (one dedicated to conventional use, and the other dedicated to disposal of biohazardous fluids and initial decontamination of equipment exposed to biohazardous fluids). A high quality water purifier has been installed. Basic utilities (gas, compressed air) are available in a large fume hood used for various chemical procedures. The lab is equipped with instrumentation for transducer characterization/exposimetry, apparatus for sonication of in vitro and in vivo biological specimens, a dual-beam spectrophotometer, a small drying oven, and is amenable to general chemistry operations (preparation of gel phantoms, etc.). See Figure 2 for photographs during and following the renovation.



Mid-way through renovation



Completed renovation

Figure 2. OFC Room 112, the CIMU Ultrasound Biophysics Lab

## OFC Room 114: Tissue Culture/Media Preparation Lab


This laboratory is composed of a larger outer room, and two small inner rooms. The outer room and one of the two inner rooms have been renovated; the second inner room was left unaltered since its primary purpose is to storage voluminous tissue culture supplies and other consumables. Photographs are shown in Figure 3. The second small inner room, which required lead-liner removal, was transformed into a dedicated tissue culture facility. Installed features include wall-mounted cabinetry for sterile supply storage, a small sink and base cabinet and a drawer unit with disinfectable work surface. The room houses a laminar flow hood for aseptic transfers, a CO<sub>2</sub> incubator, a Coulter Z1 particle counter, a clinical centrifuge, vacuum pump, and water bath. The renovated outer room contains a central island work bench, a sink and base cabinet, and several wall-mounted cabinets for glassware storage. The principal purpose of this room is for media preparation, but it was also designed to accommodate in vivo studies of small animals. Installed equipment includes an old but serviceable fume hood (existing), an ice machine, and a small autoclave for sterilization of tissue culture supplies and the degassing of water for use in ultrasound tanks. Appliances or mobile equipment/supply items include a refrigerator (media and/or biological specimen storage), balances, pH meters, chemical storage cabinet and salts, etc.


## OFC Room 125: CIMU In Vitro/Ex Vivo Studies Laboratory

This room was also completely renovated, having had only exposed concrete walls and floors and primary use as a storage area – see Figure 4. This room is located in the western wing of the building, near the CIMU ultrasound patient studies and systems development laboratories. Room 125 now is used by all CIMU staff and students to conduct bench top, in vitro experiments, involving tissue characterization, tissue-mimicking phantom production, and transducer fabrication. It also houses a wall-mounted water distiller that is heavily used by CIMU staff and students working in this far wing of the building. A small refrigerator provides chemical storage.

#### OFC Room 102M: CIMU Ultrasonic Characterization Laboratory

This room required minor renovations, including installation of plumbing and a small sink, new paint, new power and communications conduit, and custom curtains for the optical studies area. A photograph of the workspaces is provided in Figure 5. The laboratory is used by all staff and students to conduct transducer radiation force, field mapping and schlierin optical measurements. *In vitro* and *ex vivo* HIFU dose response and lesion monitoring studies are also conducted in this laboratory.









Top: Ultrasonic field characterization/in vitro/ex vivo workspace

Right: Schlierin optical system space




Figure 5. OFC Room 102M: CIMU Ultrasonic Characterization Laboratory

# C. Research Efforts

As indicated above, this grant enabled the integration of many disparate facilities and technical capabilities to provide advanced acoustic research spaces for HIFU bioengineering development and graduate education. A summary the research highlights from this project is outlined below. This list of graduate student theses supported by this grant is provided in

section II.D. A complete bibliography of research papers and presentations funded by or relevant to this grant is provided in section II.F.

## C.1 Modeling Efforts

A 2-D full-wave, fully nonlinear acoustic model was developed as part of a Ph.D. student's dissertation (see <u>Curra</u> under the list of student theses in Section II.F). This type of model is considered the "gold standard," and can be used to benchmark other, particularly approximate models, and to perform the most complete theoretical investigations into the physical and biological mechanisms of HIFU therapy. Full-wave models are highly complex, and thus require both significant computing resources as well as computing time. To reduce computational intensity, efforts also were directed toward the development of an approximate field model, in particular an implementation of a nonlinear parabolic wave equation (the KZK equation). These models have been used to perform simulation studies to support HIFU transducer design and protocol development.

## C.2 Transducer Design Studies

During the course of this grant significant progress was made toward developing a highly automated transducer characterization capability with dedicated instrumentation and facilities for this purpose. New capabilities include acoustic field mapping by scanned hydrophones, acoustic power efficiency measurement, electric and acoustic impedance measurement, acoustic and material parameter measurement, medical ultrasound imaging and RF data collection, optical methods of acoustic field mapping (Schlieren apparatus), and for each system a convenient software interface and user manual. Specialized fixtures and instruments were also developed to measure HIFU dose response *in vitro*, which forms an essential component of in the verification and development of reliable and realistic computer models for HIFU simulation and treatment planning. Single element HIFU transducers were refined with liquid, gel, and solid coupling media and other improvements to be more powerful, convenient, and robust. These various technological enhancements have enabled further development of HIFU arrays and image-guided ultrasound systems for greater flexibility and control in clinical HIFU treatment protocol.

## C.3 HIFU Monitoring Efforts

Efforts in developing HIFU monitoring techniques were focused on two efforts. The first component focused on using B-mode diagnostic imaging to monitor lesion evolution. This type of monitoring is possible because lesions formed by HIFU are hyperechoic for several seconds after treatment; CIMU studies using overpressure indicate mild cavitation is responsible for the echogenicity in the focal region. At low intensity levels the HIFU focus can be observed using interleaved B-mode imaging for a brief time (1 second or less) without causing permanent damage (i.e. no lesions were found in acute studies at such exposures). This feature enabled targeting with low HIFU exposures before applying HIFU at full treatment levels. Then, by interleaving HIFU therapy with diagnostic ultrasound interrogation, the focal region can be imaged and monitored during therapy application. This effort represents a major thrust of CIMU Image Guided Therapy (IGT) development.

The second monitoring component focused on developing new signal processing algorithms using the raw Radio Frequency (RF) ultrasound backscatter data from a diagnostic ultrasound machine, in this case the HDI-1000 developed by ATL, Inc., in Bothell, WA. Significant treatment information, including tissue motion and tissue temperature, can be obtained from the RF ultrasound data. This research effort was initiated during the latter part of the performance period and is being completed under other grants.

## C.4 HIFU Bioeffects Studies

In order to understand more fully how tissue properties change with HIFU treatment, studies were conducted in which the bulk properties of ultrasound attenuation and sound speed were measured before and after HIFU treatment. These demonstrated that ultrasound attenuation increases after HIFU treatment, whereas the speed sound remains statistically the same. These results are useful in developing treatment protocols and optimizing parameters for image-guided HIFU systems.

Instrumentation was developed to study quantitative in vitro dose response and to use in careful comparison with numerical simulations of HIFU therapy. In vitro dose response studies were initiated during the latter part of this grant and have continued under other funding vehicles. Of particular interest is investigating the threshold below which results a purely thermal effect and above which results in thermal and mechanical effects, in particular acoustic and/or vaporous cavitation.

HIFU-induced biological mechanisms leading to hemostasis in solid organs and in vascular wounds were investigated, and this effort has continued via funding from other projects. Research was directed specifically at understanding how the tissue emulsion produced during surgical treatment in intraoperative studies is formed and why it aids hemostasis.

#### C.5 Contrast Agent Studies

These studies occurred during the early part of the grant and focused on contrast agent microbubble behavior in an ultrasound field. Bubble destruction thresholds, inertial cavitation thresholds, and bubble dissolution rates were studied for a variety of contrast agents. This effort has evolved into a major area of study which is now funded by the National Institutes of Health.

# D. Degrees and Fellowships Conferred

The following individuals earned degrees through funding provided in whole or in part by this project

• Francesco Curra, Ph.D., 2001, dissertation title "Medical Ultrasound Algorithm for Noninvasive High Intensity Ultrasound Applications."

- Jonathan Yuen, M.S., 2001, thesis tittle "Characterization of lesion formation in a tissue- mimicking phantom for focused ultrasound surgery."
- Sandra Poliachik, Ph.D., 2002, dissertation title "An Investigation of the Mechanisms of High Intensity Focused Ultrasound Induced Platelet Activity."
- Tyrone Porter, Ph.D., 2003, dissertation title, "An investigation of the synergy between ultrasound and membrane-disruptive polymers and its effect on cell membranes."

The following individual's Post-Doctoral Fellowship was funded in part by this project:

• Dr. Cyril Lafon.

The following student is continuing research efforts that were initiated under this project:

 Ajay Anand, Ph.D. candidate, dissertation title "A Model-Based Noninvasive Temperature Estimation Technique for HIFU Therapy Monitoring Using Backscattered Ultrasound."

# E. Intellectual Property

One invention disclosure was filed at the University of Washington: 2031-3433 Medusa Version 1.0. The form DD 882, Report of Inventions and Disclosures, will be forward separately from the UW Office of Technology Licensing.

#### F. Publications

#### Refereed Journal Articles

Refereed Articles - 2000

Brayman AA (Section Leader), Delecki D, Wible J, Wu J, Abramowicz JS, Meltzer RS, and Porter TR. Mechanical bioeffects in the presence of gas-carrier ultrasound contrast agents. In: Fowlkes JB, Holland CK, eds., *Mechanical Bioeffects From Diagnostic Ultrasound: Consensus Statements*. J Ultrasound Med 2000; 19(2): 120-142.

Chang PP, Chen WS, Mourad PD, Poliachik SL and Crum LA. Thresholds for inertial cavitation in Albunex suspensions under pulsed ultrasound conditions. IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control 2000; 48(1):161-170.

Curra FP, Mourad PD, Khokhlova VA, Cleveland RO, and Crum LA. Numerical simulations of heating patterns and tissue temperature response due to high intensity focused ultrasound fields. IEEE Trans UFFC 2000; 47(4):1077-1089.

Dunmire B, Beach KW, Labs K, et al. Cross-beam vector Doppler ultrasound for angle-independent velocity measurements. Ultrasound Med Biol 2000; 26(8):1213-1235.

Kwok C, Mourad P, Crum L, and Ratner B. Surface modification of polymeric slab surfaces with self-assembled monolayer and is characterization with multi-surface-analytical techniques. Biomacromolecules 2000; 1(1), pp. 139-148.

Miller MW, Sherman TS, and Brayman AA. Comparative sensitivity of human and bovine Erthrocytes to sonolysis by 1 MHz ultrasound. Ultrasound Med Biol 2000; 26:1317-1326.

Refereed Articles - 2001

Bailey MR, Couret LN, Sapozhnikov OA, Khokhlova VA, ter Haar G, Vaezy S, Shi X, Martin R, and Crum LA. Use of overpressure to assess the role of bubbles in focused ultrasound lesion shape in vitro. Ultrasound Med Biol 2001 May; 27(5): 695-708.

Chang PP, Chen WS, Mourad PD, Poliachik SL, and Crum LA. Thresholds for inertial cavitation in albunex suspensions under pulsed ultrasound conditions. IEEE Trans Ultrason Ferroelectr Freq Control 2001; 48(1): 161-70.

Lafon C, Kaczkowski PJ, Vaezy S, Noble M, and Sapozhnikov OA, "Development and characterization of an innovative synthetic tissue-mimicking material for high intensity focused ultrasound (HIFU) exposures," IEEE Ultrasonics Symposium, 2, 1295-1298 (2001).

Mourad PD, Lazar DA, Curra FP, Mohr BC, Andrus KC, Avellino AM, McNutt LD, Crum LA, and Kliot M. Ultrasound accelerates functional recovery after peripheral nerve damage. Neurosurgery 2001; 48(5): 1136-41.

Mourad PD, Murthy N, Porter TM, Poliachik SL, Crum LK, Hoffman AS, Stayton PS. Focused ultrasound and poly(2-ethylacrylic acid) act synergistically to disrupt lipid bilayers in vitro. Macromolecules 2001; 34(8): 2400-2401.

Poliachik SL, Chandler WL, Mourad PD, Ollos RJ, Crum LA. Activation, aggregation and adhesion of platelets exposed to high-intensity focused ultrasound. Utrasound Med Bio 2001; 27(11):1567-1576.

Porter T, Crum L, Stayton P, Hoffman A, "Sonoporation of Erythrocytes by Combination of Synthetic Polymer and Ultrasound," In: Proc. 17<sup>th</sup> International Congress on Acoustics (Rome, Italy, Sept 2-7, 2001).

Shi X, Martin RW, Vaezy S, Kaczkowski P, and Crum LA. Color Doppler detection of acoustic streaming in a hematoma model. Ultrasound Med Biol 2001; 27(9): 1255-64.

Vaezy S, Martin RW, and Crum LA. Acoustic Surgery. Physics World 2001; 14(8): 35-39.

Vaezy S, Martin RW, and Crum LA. High intensity focused ultrasound: a method of hemostasis. Echocardiography 2001; 18(4): 309-15.

Vaezy S, Shi XG, Martin RW, Chi E, Nelson PI, Bailey MR, and Crum LA. Real-time visualization of high-intensity focused ultrasound treatment using ultrasound imaging. Ultrasound Med Biol 2001;27(1): 33-42.

#### Refereed Articles - 2002

Anand AJ, Kaczkowski PJ, Daigle RE, Huang L, Paun M, Beach KW, and Crum LA, "Using the ATL HDI-1000 ultrasound scanner to collect demodulated RF data for monitoring HIFU lesion formation," *Proc SPIE*, **5035**, 316-326 (2003).

Curra FP, Kargl SG, and Crum LA, "Parameter space investigation of optimal thermal lesion generation in noninvasive HIFU applications," in <u>Therapeutic Ultrasound, Proceedings of the 2<sup>nd</sup> International Symposium</u>, M.A. Andrew, L.A. Crum and S. Vaezy, eds (American Institute of Physics Press), pp. 275-281 (2003).

#### Refereed Articles - 2003

Anand AJ, Kaczkowski PJ, "Monitoring formation of High Intensity Focused Ultrasound (HIFU) induced lesions using backscattered ultrasound", Acoustic Research Letters Online, August 2003 (submitted).

Curra FP and Crum LA, "Therapeutic ultrasound: Surgery and drug delivery," Acoustic Science and Technology, 24, 6 (2003).

Poliachik SL, Chandler, WL, Ollos RJ, Bailey MR, and Crum, LA, "The Relation Between Cavitation and Platelet Aggregation During Exposure to High Intensity Focused Ultrasound," *Ultrasound Med Biol*, accepted, (2003).

# **Books or Book Chapters**

Andrew MA, Crum LA, and Vaezy S (editors), Proceedings of the 2nd International Symposium on Therapeutic Ultrasound, University of Washington, Seattle, Washington, 2003.

Crum, L.A., K. Beach, S. Carter, W. Chandler, F. P. Curra, P. Kaczkowski, G. Keilman, V. Khokhlova, R. Martin, P. D. Mourad, and S. Vaezy (2000) Acoustic Hemostasis. In: W. Lauterborn and T. Kurz, (eds.), "Nonlinear Acoustics at the Turn of the Millennium," Am. Inst. Of Physics, ISNA 15, (New York), pp. 13-22.

Crum LA, Bailey M, Carter S, Curra F, Kaczkowski P, Kargl S, "Image-guided acoustic hemostasis," in <u>New Acoustics: Selected Topics</u>, C. Ranz-Guerra and J. A. Gallego-Juarez, eds., (Consejo Superior de Investigaciones Cientificas, Madrid), pp 26-36 (2002).

D. A. Lazar, F. P. Curra, B. C. Mohr, L. D. McNutt, M. Kliot and Mourad, P. D., "Acceleration of recovery after injury to the peripheral nervous system via ultrasound and other therapeutic modalities", In <u>Neurosurgery Clinics of North America</u> -Peripheral Nerve

<u>Issues: Controversies and Evolving Treatments-</u>, M. Kliot editor, W. B. Saunders Publishing House, April 2001.

Vaezy, S. Martin RW, Kaczkowski P, Keilman G, Goldman B, Yaziji H, Carter S, Caps M, Crum LA: Use of High Intensity Focused Ultrasound to Control Bleeding. Yearbook of Vascular Surgery 2000.

Vaezy S, Andrew M, Kaczkowski P, and Crum L. Image-guided acoustic therapy. Annu Rev Biomed Eng 2001; 3: 375-90.

## **Technical Reports**

Mourad, P.D. and S.G. Kargl (2000) Acoustic Properties of Fluid-Saturated Blood Clots. APL-UW TR 2003.

#### **Student Theses**

Curra, F, "Medical Ultrasound Algorithm for Noninvasive High Intensity Ultrasound Applications" - Ph.D. Dissertation, University of Washington, August, 2001.

Poliachik, S. "An Investigation of the Mechanisms of High Intensity Focused Ultrasound Induced Platelet Activity," Ph.D. Dissertation, University of Washington, 2002.

Porter, T., "An investigation of the synergy between ultrasound and membrane-disruptive polymers and its effect on cell membranes"

Yuen, J, "Characterization of lesion formation in a tissue-mimicking phantom for focused ultrasound surgery," M.S. Thesis, University of Washington, December, 2001.

#### Abstracts, Posters & Presentations

Abstracts & Presentations - 2000

Crum L, Bailey M, Carter S, Curra F, Kaczkowski P, Kargl S, Martin R, Mourad P, and Vaezy S, "Image-guided Acoustic Hemostasis." Symposium on Architectural Acoustics, II Ibero-American Congress of Acoustics, XXXI National Congress of Acoustics - TECNIACUSTICA 2000-, II Ibero-American Meeting on Ultrasonic and II Iberian Congress of Acoustics, Madrid, Spain, 16-20 October 2000

Curra FP, Mourad PD, Khokhlova VA, and Crum LA, "3D full wave ultrasonic field and temperature simulations in biological tissue containing a blood vessel." 139th Meeting of the Acoustical Society of America, Atlanta, GA, May 30-June 3, 2000

Curra FP, Mourad PD, Kargl SG, and Crum LA, "Theorectical predictions of ultrasound fields, temperature response, and lesion dynamics in biological tissue for the purpose of noninvasive disease treatment," *J. Acoust. Soc. Am*, 108, 2546 (2000).

Lafon C, Bailey M, Couret L, Kaczkowski P, Sapozhnikov O, Brayman A, Crum L. Real-time observation of inception and growth of HIFU-induced tissue lesions. J. of Acous. Soc. of Am., Newport Beach, USA, 2000;108:2546.

Martin R, Vaezy S, Shi X, Kaczkowski P, paun M, Beach K, and Crum L (2000) "Real-Time Visualization of Therapeutic Ultrasound: Applied to Acoustic Hemostasis," 9<sup>th</sup> Congress of the World Federation for Ultrasound in Medicine and Biology, Florence Italy.

Mourad, P.D., D.A. Lazar, F.P. Curra, B. Mohr, K.C. Andrus, A.M. Avellino, L.D. McNutt, L.A. Crum and M. Kliot, *Ultrasound accelerates functional recovery after peripheral nerve damage*, 9th International Meeting of the World Federation of Ultrasound in Medicine and Biology, Florence, Italy, 6-10 May, 2000.

Poliachik SL, Chandler WL, Mourad PD, Ollos RJ, Crum LA. Effect of high-intensity focused ultrasound on platelet aggregation, activation and adhesion. Institute of Cancer Research ,Joint Department of Physics, Royal Marsden Hospital, Surrey England, 26 April 2000.

Poliachik, SL, Chandler, WL, Mourad, PD, Ollos, RJ, Crum, LA. Effect of high-intensity focused ultrasound on platelet aggregation, activation and adhesion. 9th Congress of the World Federation for Ultrasound in Medicine and Biology, Florence, Italy, 6-10 May 2000.

Poliachik SL, Chandler WL, Mourad PD, Ollos RJ, Crum LA. Effect of high-intensity focused ultrasound on platelet aggregation, activation and adhesion. 139th Meeting Acoustical Society of America, Atlanta, GA, 31 May 2000.

Poliachik SL, Chandler WL, Mourad PD, Ollos RJ, Crum LA. Activation, Aggregation and Adhesion of Platelets Exposed to High Intensity Focused Ultrasound. 2000 IEEE Ultrasonics Symposium Proceedings; 2000; Vol. 2 pp. 1433-1436.

Poliachik, SL, Chandler, WL, Mourad, PD, Ollos, RJ, Crum, LA. Platelet activity as a result of exposure to high intensity focused ultrasound. 140th Meeting Acoustical Society of America, Newport Beach, CA, 6 December 2000.

Porter, TM, Nickerson J, Crum LA, Black FE, Murthy N, Stayton PS, and Hoffman AS, "Influence of chemical composition of membrane-disrupting polymers on relative cavitation activity and hemolysis," *J Acoust Soc Am*, **108** (5), Pt. 2, 2547 (2000).

Shi X, Martin R, Vaezy S, and Crum L (2000) Experimental Investigation and Finite Element Simulation of Streaming in Blood in Cylindrical Models, IEEE International Ultrasonics Symposium.

Vaezy S, Martin R, Shi X, Paun M, Beach K, Kaczkowski P, Keilman G, Carter S, Bailey M, and Crum L (2000) Hemostasis of Catheter-Induced Femoral Artery Injuries using Image-Guided Transcutaneous High Intensity Focused Ultrasound, Annual Biomedical Engineering Society Meeting, Seattle, WA.

#### Abstracts & Presentations - 2001

Bailey MR, Vaezy S, Yuen JC, Anand A, Miller NA, Kaczkowski PJ, and Crum LA. Bubbles and acoustic image-guided high intensity focused ultrasound. 142<sup>nd</sup> Meeting Acoustical Society of America, 3-7 December 2001, Fort Lauderdale, FL, J Acoust Soc Am 2001; 110: 2643.

Chen W, and Matula T. A light-scattering technique for investigating ultrasound contrast agents. Session 1D-5, p. 144. 2001 IEEE International Ultrasonics Symposium, 7-10 October 2001, Atlanta, GA

Crum LA. Acoustic hemostasis. Session 2H-3, p. 286. 2001 IEEE International Ultrasonics Symposium, 7-10 October 2001, Atlanta, GA

Curra FP, Kargl SG, Lafon C, and Crum LA, "Theoretical predictions and experimental results for non-invasive disease treatment via high intensity focused ultrasound: a comparative study," 17th International Congress on Acoustics 2001, Rome, Italy, Biomedicine: Therapeutic Ultrasound, 4, 20-21 (2001).

Curra FP, Kargl SG, Lafon C, and Crum LA, "Thermal lesions produced by High Intensity Focused Ultrasound: Theoretical predictions and experimental results," Proc. of First International Workshop on the Application of High Intensity Focused Ultrasound (HIFU) in Medicine 2001, Chongqing, China, 29-31.

Curra FP, Kargl SG, Lafon C, and Crum LA, "High intensity focused ultrasound for non-invasive disease treatment: Theoretical predictions and experimental results," 141<sup>st</sup> Meeting of the Acoustical Society of America, Chicago, IL, June 4-8, 2001, J Acoust Soc Am, 109 (5), p. 2457 (2001).

Dunmire B and Beach KW (2001). A brief history of vector Doppler. SPIEs International Symposium Medical Imaging 2001, San Diego, CA, USA.

Dunmire B, Beach KW, Labs K-H, Paun M, Tschoeppel M (2001). Two-dimensional velocity map of a normal femoral bifurcation, and its amplications for conventional pulsed Doppler ultrasound. SPIEs International Symposium Medical Imaging 2001, San Diego, CA, USA.

Dunmire B, Pagel G, Beach KW, Labs K-H (2001). Post stenotic flow distrurbances in a steady flow model. SPIEs International Symposium Medical Imaging 2001, San Diego, CA, USA.

Kaczkowski PJ, Vaezy S, Martin RW, and Crum LA. A multi-channel high-intensity focused ultrasound system for image-guided therapy. J Acoust Soc Am 2001; 110: 2614. 142<sup>nd</sup> Meeting Acoustical Society of America, 3-7 December 2001, Fort Lauderdale, FL

Lafon C, Vaezy S, Noble ML, Kaczkowski PJ, Martin RW, and Crum LA, "A new synthetic tissue-mimicking phantom for high intensity focused ultrasound," 17th International Congress on Acoustics, Biomedicine: Therapeutic Ultrasound, 7, 32-33 (2001).

Lafon C, Vaezy S, Noble ML, Kaczkowski PJ, Martin RW, and Crum LA. A new synthetic tissue-mimicking phantom for high intensity focused ultrasound. 17<sup>th</sup> International Congress on Acoustics 2001. Volume 4. Biomedicine. Therapeutic Ultrasound. Pp. 32-33. 17<sup>th</sup> International Congress on Acoustics, 2-7 September 2001, Rome, Italy

Lafon C, Kaczkowski PJ, Vaezy S, Sapozhnikov OA, and Noble ML. Development and characterization of an innovative synthetic tissue-mimicking material for high intensity focused ultrasound (HIFU) exposures. Session P1J-2. P. 303. 2001 IEEE International Ultrasonics Symposium,7-10 October 2001, Atlanta, GA

Lafon C, Sapozhnikov OA, Kaczkowski PJ, Vaezy S, Noble ML, and Crum LA. An innovative synthetic tissue-mimicking material for high-intensity focused ultrasound. J Acoust Soc Am 2001; 110: 2613. 142<sup>nd</sup> Meeting Acoustical Society of America, 3-7 December 2001, Fort Lauderdale, FL

Matula TJ, and Chen WS. A light-scattering technique for investigating ultrasound contrast agents. 17<sup>th</sup> International Congress on Acoustics 2001. Volume 4. Biomedicine. Acoustics in Medicine. Pp. 4-5. 17<sup>th</sup> International Congress on Acoustics, 2-7 September 2001, Rome, Italy

Poliachik SL, Chandler WL, Ollos RJ, and Crum LA. Role of high-intensity focused ultrasound induced cavitation on platelet aggregation. 17<sup>th</sup> International Congress on Acoustics 2001. Volume 4. Biomedicine. Medical Ultrasound Bioeffects. Pp. 20-21. 17<sup>th</sup> International Congress on Acoustics, 2-7 September 2001, Rome, Italy

Shi X, Martin RW, Vaezy S, and Kaczkowski PJ. Color Doppler imaging of acoustic streaming for hematoma diagnosis. Session 1G-6. P. 261. 2001 IEEE International Ultrasonics Symposium,7-10 October 2001, Atlanta, GA

Vaezy S. High intensity focused ultrasound for therapy in medicine. 17<sup>th</sup> International Congress on Acoustics 2001. Volume 4. Biomedicine. Therapeutic Ultrasound. Pp. 2-3. 17<sup>th</sup> International Congress on Acoustics, 2-7 September 2001, Rome, Italy

Vaezy S, Martin RW, and Crum LA. High intensity focused ultrasound for arrest of bleeding. J Acoust Soc Am 2001; 110: 2643. 142<sup>nd</sup> Meeting Acoustical Society of America, 3-7 December 2001, Fort Lauderdale, FL

#### Abstracts & Presentations - 2002

Anand A, Huang L, Kaczkowski P, Daigle R, Crum L, "Using ATL HDI-1000 Ultrasound Scanner for Tissue Elasticity Imaging," Proceedings of the First International Conference on

the Ultrasonic Measurement and Imaging of Tissue Elasticity, Niagara Falls, Ontario, Canada (Oct. 2002).

Curra FP, Kargl SG, and Crum LA, "3-D Full Wave Ultrasonic Field and Temperature Simulations in Inhomogeneous Biological Tissue," Proceedings of the 16th International Symposium on Nonlinear Acoustics 2002, Moscow, Russia, 19-23.

Curra FP, Kargl SG, and Crum LA, "Parameter space investigation of optimal thermal lesion generation in noninvasive HIFU applications," in <u>Therapeutic Ultrasound</u>, <u>Proceedings of the 2<sup>nd</sup> International Symposium</u>, M.A. Andrew, L.A. Crum and S. Vaezy, eds (American Institute of Physics Press), pp. 275-281 (2003).

Curra FP, Kargl SG, Lafon C, and Crum LA, "Theoretical predictions and experimental results for non-invasive disease treatment via High-Intensity Focused Ultrasound: a comparative study", Proceedings of the 17th International Congress on Acoustics, Rome, Italy, 2-7 September 2001.

Abstracts & Presentations - 2003

Anand AJ, Kaczkowski PJ, "Monitoring evolution of HIFU-induced lesions with backscattered ultrasound", 145th Meeting of the Acoustical Society of America, Nashville, TN, April, 2003, J. Acoust. Soc. Am., Vol. 113, No. 4, Pt. 2, p 2310 (2003).

Anand AJ, Kaczkowski PJ, "Ultrasound RF-signal analysis of HIFU-induced lesions", Abstracts of 28<sup>th</sup> Ultrasonic Imaging and Tissue Characterization Symposium, Arlington, VA published in Ultrasonic Imaging, Vol. 25 (1), Jan 2003.